

Quantitative evaluation of regional precipitation forecasts using multi-dimensional remote sensing observations (QUEST)

Susanne Crewell¹, and the QUEST Team


crewell@meteo.uni-koeln.de <http://gop.meteo.uni-koeln.de>

¹ IGMK: Universität zu Köln, Institut für Geophysik und Meteorologie, Kerpener Str. 13, 50923 Köln




Objectives

Precipitation is the final atmospheric process of the hydrological cycle. Consequently quantitative precipitation forecasts (QPF) can only be successful, if a model represents all processes of this cycle accurately. The project "Quantitative evaluation of regional precipitation forecasts using multi-dimensional remote sensing observations" (QUEST) aims at a complete analysis of the modeled hydrological cycle in order to identify the reasons of QPF deficiencies and to give distinct advices for model improvement.



Case Studies



In order to demonstrate the QUEST tools two case studies (stratiform 3 August 2006 and convective 28 August 2006) were performed [Pfeifer et al., 2009]. In order to optimally exploit satellite observations forward operators were applied to simulate synthetic brightness temperatures (BT) from the model output.



- MSG Infrared observations indicate widespread system with high clouds over Germany whose cloud top height is underestimated
- AMSU Microwave observations indicate underestimation of amount of highly scattering frozen precipitation

MSG Diurnal Cycle

- Assimilation of radiosondes with dry bias (noon time) causes jump in bias for runs started at 12, 15 and 18 UTC; e.g. 6.2 and 10.8 μm TB are too high because of lower water vapor and lesser high clouds.
- Diurnal cycle of 10.8 μm BT is a superposition of diurnal surface warming and diurnal cloudiness. COSMO-EU shows a phase shift of ~1h.
- Cloud mask is overestimated in the afternoon and underestimated during night. THS and TSS perform slightly better during early morning hours.



Stefan Stapelberg, FU Berlin



Publications:

Crewell, S., M. Mech, T. Reinhardt, C. Selbach, H.-D. Betz, E. Brocard, G. Dick, E. O'Connor, J. Fischer, T. Hanisch, T. Hauf, A. Hünerbein, L. Delobbe, A. Mathes, G. Peters, H. Wernli, M. Wiegner and V. Wulfmeyer, 2008: The General Observation Period 2007 within the Priority Programm on Quantitative Precipitation Forecasting. Concept and first results. *Meteorol. Z.* 17(6), 849-866.
 Pfeifer, M., W. Yen, M. Hagen, G. Craig, T. Reinhardt, M. Mech, S. Crewell, A. Hünerbein, J. Fischer, M. Schröder, and M. Baldauf, 2008: Validating precipitation forecasts using sensor synergy: The case study approach. *Meteorol. Z.* submitted.

Long-term Validation using GOP


During the General Observation Period (GOP, Crewell et al. 2008) started in the beginning of 2007 a comprehensive data set of observations and corresponding model forecasts by the operational forecasts of DWD's COSMO-DE ($\Delta x \approx 2.8$ km) and COSMO-EU ($\Delta x \approx 7$ km) models is gathered:

- Satellites:**
- SEVIRI on MSG
 - AMSU on NOAA/Metop
 - Cloudsat
- Other sensors:**
- Radiosonde
 - Cloudnet
 - Micro Rain Radar
 - GPS
 - Celiometer
 - Weather Radar




Time series of integrated water vapor (IWV, top) observed by GPS network, cloud base height observed by celiometer network (middle) and precipitation observed by Ranie1 (gauge only) and Ranie2 (gauge/radar) together with closest (blue) and +12 h (magenta) forecast for COSMO-DE (solid) and COSMO-EU (dotted).

IWV & Cloud Base Diurnal cycle



Time series of COSMO-De 12h forecast CLB RMSE as a function of day time (color coded).

Suraj Polade, UHamburg



- Data assimilation has strong influence on diurnal cycle of closest forecasts.
- After 12h forecast COSMO-DE has developed a reasonable diurnal IWV cycle (1.0 vs 0.8 kgm^{-2} observed).
- Diurnal cycle in cloud base is predicted too weakly while seasonal cycle is predicted too strongly.
- Cloud base is predicted worst during night in summer revealing problems in the parametrization of stable boundary layer.

Dependence on Weather Type

Tom Akkermans, KU Leuven

- Classification into weather types (8 wind directional and 2 vorticity classes; cyclonic C + anticyclonic AC) after Jenkinson and Collison (1977) based on COSMO-EU output on 3 hourly base for whole Germany at 850hPa.
- During southerly flows to too humid conditions especially in South Germany connected with too low cloud bases occur.
- Precipitation is underestimated during convective situations (stability diagnosed from radiosondes).

